

## **Total Hip Replacement - Anterior First Post Operative Visit**

**Sutures:** Anterior hip incisions are typically closed with dermabond moncryl (glue). However, in certain cases staples will be used for closure, make sure to examine incision site. If so, staples will generally be removed at this visit if the wound is healing well. In some circumstances, staples will be left in for an additional week. After staples are removed you may continue to shower daily. Small steri-strips on the wound should be removed in 1 week. If any fall off, the steri-stripes do not need to be replaced before 1 week is over. When showering, let soap and water run over the incision (do not scrub). Do not submerge the incision in a pool or bath tub for another 3 weeks. The incision should no longer have any drainage after the staples are removed; a bandage is NO longer necessary. If any wound drainage develops, please call us immediately.

**Pain Medication:** Some patients require some pain medication for up to 6 weeks after surgery, particularly for physical therapy and perhaps to help with sleeping. We will be happy to provide refills during this time if necessary. Now is an appropriate time to start decreasing the amount of medicine you are taking. Tylenol can be substituted for the narcotic. Do not exceed 3000mg of Tylenol in 24 hours and remember that the pain medication may also contain Tylenol – this must be considered when calculating the total dose. Do not use anti-inflammatory medications (Motrin, Advil, Aleve, etc.) other than Celebrex while taking blood thinners unless approved by our team. Medication refills must be requested during business hours (8am-4:30pm) and at least 24-48 hours before the prescription runs out. The best way to request a refill for medication is through Mychart or via phone if needed at (410) 674-1641.

**Physical Activity:** Unless you have been given special precautions, you may bear weight as tolerated and should continue with your home exercise program two times a day. Physical therapy is an important component of your recovery, and should be continued, usually 2 times a week. You may start to use cane, and then no assistive devices, under the supervision of your physical therapist. You may sleep on your side, with a pillow between your knees. DO NOT sleep with any pillows or rolls under the knee – this will encourage stiffness.

**DVT prevention (Aspirin):** Continue Aspirin 325mg two times a day for 6 weeks after surgery. TED stockings may be discontinued 4 weeks after surgery, although you may wear them longer if you like, particularly if you have some leg swelling. For patients requiring additional forms of anti-coagulation, specific instructions will be provided for that medication. Please call if there are any questions. In general, if you are on eliquis after surgery, this will be stopped 4 weeks after the surgery, and then you should begin taking ecotrin 81mg twice a day for 2 additional weeks

**General Notes:** Many patients will feel some fatigue, have trouble sleeping, and report some leg swelling after joint replacement surgery. This is normal, and should resolve over the next few weeks. If you have problems or concerns at any time, please do not hesitate to contact us at the office. For emergencies after hours, a physician is available 24 hours a day, 7 days a week can be reached at (410)-268-8862.